



MISSISSIPPI DEPARTMENT OF
ENVIRONMENTAL QUALITY
OFFICE OF GEOLOGY
OPEN - FILE REPORT 187

GEOLOGIC MAP of the HESTERVILLE QUADRANGLE Attala and Carroll Counties, Mississippi



Geology by David E. Thompson, RPG



2004

DESCRIPTION OF MAP UNITS


QUATERNARY	PLEISTOCENE	HOLOCENE		
			Symbol	Description
QUATERNARY	PLEISTOCENE	HOLOCENE	Qal	ALLUVIUM Sand, flood plain sands and silts.
			Tk	LOESS Silt, buff to tan, pale yellow, red, or gray, sandy to clayey, quartzose, feldspathic. Unweathered loess is typically calcareous with dolomite and calcite; however, loess in this quadrangle is highly weathered, leached/noncalcareous, very clayey, and has been referred to as a brown or yellow loam. Loess is an eolian deposit derived from glacial outwash. Loess deposits blanket the pre-loess topography of the quadrangle area, with greater quantities developed along ridge crests than in valleys, creating local variation in thickness. The thickness in the quadrangle is estimated at 2 to 7 feet; loess is present east of the mapped area, but coverage is discontinuous and the average thickness is less than 5 feet. In places, weathered loess contains secondary deposits of small calcareous concretions (caliche, loess dolls).
			Tk	KOSCIUSKO FORMATION Sand, gray to light olive gray, weathers reddish orange to pale yellowish brown, very fine- to very coarse-grained, quartzose, micaceous; interbedded to interlaminated with silt and clay, light olive gray to brownish gray, carbonaceous to lignitic; especially argillaceous in upper third of the formation. Locally, the basal Kosciusko contains layers of quartzite, siliceous siltstone and sandstone as thick as 5 feet, often occurring as large boulders along hill tops and slopes. Unconformity at base. The thickness is estimated to be 300 feet; however, only the lower 140 feet or so are exposed in the southern portion of the quadrangle. Constitutes the Sparta Aquifer.
TERTIARY	EOCENE	CLAIBORNE GROUP	Twn-Tz	ZILPHA and WINONA FORMATIONS Zilpha - Clay, gray to brownish black, carbonaceous to lignitic, weathers light gray to reddish pink to white, massive and homogeneous or interbedded to interlaminated with silt and sand, gray to light olive gray, quartzose, micaceous, carbonaceous, locally glauconitic; concretionary siderite and limonite; near surface exposures may exhibit jointing with selenite or limonite infilling. The thickness is variable from a few feet to 60 feet. Winona - Sand, gray to green, weathers very light gray to reddish orange or dark red, fine- to coarse-grained, quartzose, micaceous, typically glauconitic to very glauconitic, carbonaceous, silty, locally fossiliferous with thin marine shell beds and prints. Surface exposures commonly weather to distinctive contorted, concretionary, limonitic sandstone and sandy ironstone, concretionary siderite, especially near top. Approximately 60 feet thick. The total thickness of the Zilpha/Winona interval is approximately 120 feet.
			Tbc	TALLAHATTA FORMATION Basic City Member Sand, gray to very light gray, weathers pale yellowish orange to reddish orange, very fine- to medium-grained, unconsolidated, massive to cross-bedded, quartzose, micaceous, carbonaceous, pyritic; also greenish yellow to buff, fine-grained, semi-consolidated, siliceous, glauconitic, and silty; interbedded to interlaminated with clay, silt, claystone, and quartzitic siltstone and sandstone, olive gray to brownish gray, weathers yellowish gray to very light gray or white, carbonaceous with leaf and plant impressions, fucoidal structures are common, near surface exposures may exhibit jointing with limonite infilling; claystones typically weather to lightweight and brittle rock with a subconchoidal fracture. Unconsolidated sands in the upper 30 to 60 feet are equivalent to the Neshoba Sand interval. The total thickness is approximately 220 feet; however, only the upper 90 feet or so are exposed in the northeastern portion of the quadrangle.



GEOLOGIC MAP
HESTERVILLE QUADRANGLE
Attala and Carroll Counties,
Mississippi



Copyright © 2004 Mississippi Department of Environmental Quality, Office of Geology



Geology field checked in 2003 using the 1964 U.S. Geological Survey 7.5-minute topographic quadrangle, 1927 North American datum, contour interval 10 feet. Mississippi Transverse Mercator projection, 1983 North American datum, GR580 spheroid, 1000-meter Universal Transverse Mercator grid ticks, zone 16; 1983 datum shown in red, 1927 datum shown in blue.

Sources: Road and water features, USGS Digital Line Graph data, 1:100,000 scale. Public Land Survey System and contours, Mississippi Automated Resource Information System (MARIS), 1:24,000 scale.

Geographic Information System by Daniel W. Morse. MDEQ does not warrant the accuracy or completeness of the source data. maps are only a guide to current understanding and do not eliminate the need for detailed investigations of specific sit

This map was produced by the Mississippi Office of Geology in cooperation with the U.S. Geological Survey, National Geologic Mapping Program, under STATEMAP grant #03F1QA0073.