

MISSISSIPPI DEPARTMENT OF ENVIRONMENTAL QUALITY
 OFFICE OF GEOLOGY
 OPEN-FILE REPORT 198
GEOLOGIC MAP
 of the
BAILEY LAKE QUADRANGLE
 Carroll and Montgomery
 Counties, Mississippi



Geology by David E. Thompson, RPG

2005

DESCRIPTION OF MAP UNITS



- | | | | | |
|------------|-----------------|-------------------------------------|--------|--|
| QUATERNARY | HOLOCENE | ALLUVIUM | Qal | Sand, flood plain sands and silts. |
| | PLEISTOCENE | LOESS | Lo | Silt, buff to tan, pale yellow, red, or gray, sandy to clayey, quartzose, feldspathic. Unweathered loess is typically calcareous with dolomite and calcite; however, loess in this quadrangle is highly weathered, leached/noncalcareous, very clayey, and has been referred to as a brown or yellow loam. Loess is an eolian deposit derived from glacial outwash. Loess deposits blanket the pre-loess topography of the quadrangle area, with greater quantities developed along ridge crests than in valleys, creating local variation in thickness. The thickness in the quadrangle is estimated at 5 to 7 feet. In places, weathered loess contains secondary deposits of small calcareous concretions (caliche, loess dolls). |
| TERTIARY | EOCENE | KOSCIUSKO FORMATION | Tk | Sand, gray to light olive gray, weathers reddish orange to pale yellowish brown, very fine- to very coarse-grained, quartzose, micaceous; interbedded to interlaminated with silt and clay, light olive gray to brownish gray, carbonaceous to lignitic, especially argillaceous in upper third of the formation. Locally, the basal Kosciusko contains layers of quartzitic, siliceous siltstone and sandstone as thick as 5 feet, often occurring as large boulders along hill tops and slopes. Unconformity at base. The thickness is estimated to be 300 feet; however, only the lower 220 feet or so are exposed in the quadrangle. Constitutes the Sparta Aquifer. |
| | CLAIBORNE GROUP | WINONA and ZILPHA FORMATIONS | Twn-Tz | Zilpha - Clay, gray to brownish black, carbonaceous to lignitic, weathers light gray to reddish pink to white, massive and homogeneous or interbedded to interlaminated with silt and sand, gray to light olive gray, quartzose, micaceous, carbonaceous, locally glauconitic; locally fossiliferous with thin marine shell beds and prints. Surface exposures commonly weather to distinctive contorted, concretionary, limonitic sandstone and sandy ironstone; concretionary siderite, especially near top. Approximately 60 feet thick.

Winona - Sand, gray to green, weathers very light gray to reddish orange or dark red, fine- to coarse-grained, quartzose, micaceous, typically glauconitic to very glauconitic, carbonaceous, silty, locally fossiliferous with thin marine shell beds and prints. Surface exposures commonly weather to distinctive contorted, concretionary, limonitic sandstone and sandy ironstone; concretionary siderite, especially near top. Approximately 60 feet thick.

The total thickness of the Zilpha/Winona interval is approximately 120 feet; however, the unit thins to 80 feet or so locally due to apparent overlap by Kosciusko sand. |
- J - 1 Drill-hole locality and identification number



GEOLOGIC MAP
BAILEY LAKE QUADRANGLE
 Carroll and Montgomery Counties,
 Mississippi



Copyright © 2006 Mississippi Department of Environmental Quality, Office of Geology


Geology field checked in 2004 using the 1975, photorevised 1982, U.S. Geological Survey 7.5-minute topographic quadrangle, 1927 North American datum, contour interval 20 feet. Universal Transverse Mercator projection, 1983 North American datum, GRS80 spheroid, 1000-meter Universal Transverse Mercator grid ticks, zone 16; 1983 datum shown in red. January 2005, magnetic north declination in quadrangle center is 0°4' east of true north.

Sources: Road features, USGS Digital Line Graph data, 1:100,000 scale. Water features, USGS National Hydrography Dataset, 1:24,000 scale. Public Land Survey System and contours, Mississippi Automated Resource Information System (MARIS), 1:24,000 scale. Declination, National Oceanic and Atmospheric Administration (NOAA).

Geographic Information System by Daniel W. Morse, MDEQ does not warrant the accuracy or completeness of the source data. Geologic maps are only a guide to current understanding and do not eliminate the need for detailed investigations of specific sites for specific purposes.

This map was produced by the Mississippi Office of Geology in cooperation with the United States Geological Survey, National Geologic Mapping Program, under STATEMAP grant #04HQAG0031.

Structural Cross-Section of the Bailey Lake 7.5-Minute Geologic Quadrangle



Vertical Exaggeration X 15

