



GEOLOGIC MAP
SOSO QUADRANGLE
Jones, Smith, and Jasper
Counties, Mississippi



Copyright © 2011 Mississippi Department of Environmental Quality, Office of Geology

Geology field checked in 2011 using the 1974, U.S. Geological Survey 7.5-minute topographic quadrangle, 1983 North American datum, contour interval 10 feet, 1000-meter Universal Transverse Mercator grid ticks, zone 16, 1983 datum shown in red January 2011, magnetic north declination in quadrangle center is 0°48' west of true north.

Sources: The base map is derived from a Digital Raster Graphic of the USGS topographic quadrangle map, Declination, National Oceanic and Atmospheric Administration (NOAA).

Geographic Information System by Daniel W. Morse, MDEQ does not warrant the accuracy or completeness of the source data. Geologic maps are only a guide to current understanding and do not eliminate the need for detailed investigations of specific sites for specific purposes.

This map was produced by the Mississippi Office of Geology in cooperation with the United States Geological Survey, National Geologic Mapping Program, under STATEMAP grant #G10AC00294.



MISSISSIPPI DEPARTMENT OF
ENVIRONMENTAL QUALITY
OFFICE OF GEOLOGY
OPEN-FILE REPORT 244

GEOLOGIC MAP
of the
SOSO QUADRANGLE

Jones, Smith, and Jasper
Counties, Mississippi



Geology by James E. Starnes, RPG

2011

DESCRIPTION OF MAP UNITS

| | | | | | |
|-------------|-----------------|----------------|----------|-----------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| QUATERNARY | HOLOCENE | <div>Qal</div> | ALLUVIUM | Flood plain sands, silts, gravels, and clays. | |
| | | | | | |
| PLEISTOCENE | PLIO-PLISTOCENE | <div>Qtc</div> | | CITRONELLE FORMATION | Sand, yellow, orange, purple, red, pink, fine- to coarse-grained, predominantly quartzose, cross-bedded to massive; graveliferous, pea to cobble size, predominantly chert with lesser amounts of vein quartz, metaquartzite, agate, and sandstone; clay, pink to white, generally occurring as discontinuous lenses and as rip-up clasts, clasts may be boulder size. Conglomeratic ironstone ledges are common in the graveliferous sands at the base of the formation, which overlies the Catahoula and Hattiesburg Formations unconformably. |
| | | | | | |
| TERTIARY | MIocene | <div>Tha</div> | | HATTIESBURG FORMATION | Clay, green, gray, brown, weathers white to brown and contains opaline concretions in places, silty to sandy (silt commonly weathers to mottled reddish-purple and gray, dense, ferruginous concretionary masses), locally lignitic; sand, gray, pale yellow to white, fine- to coarse-grained, cross-bedded to massive, containing pea gravel in basal portion, often indurated to sandstones and siltstones at the surface, predominantly quartzose with lesser amounts of chert, metaquartzite, mica, and heavy minerals, silicified and coalified wood common; gravel, well-rounded quartz (white, yellow, brown, pink, and clear), agate (gray, yellow, white, banded, quartz druse or chalcedony), and subangular to well-rounded chert (white, gray, black). Some chert clasts are oolitic, banded, or contain marine Paleozoic fossils such as crinoids, brachiopods, bryozoans, rugose and tabulate corals, and gastropods. The base of the Hattiesburg Formation is designated at the base of a sand unit of regional extent that occurs at the approximate horizon of the base of the Fleming Formation in Louisiana and the middle-Miocene Amos Sand in Alabama. |
| | | | | | |
| CENOZOIC | OLIGOCENE | <div>Tca</div> | | CATAHOULA FORMATION | Sand, gray, pale yellow to white, fine- to coarse-grained, cross-bedded to massive with rare thinly-bedded pea gravels (gravels consist of black chert and milky quartz, are highly polished, subangular to well rounded), often indurated to sandstones at surface, predominantly quartzose with lesser amounts of chert, metaquartzite, mica, and heavy minerals, slightly glauconitic in places, silicified wood and fossil palm common; clay, green, gray, brown, weathers white to brown, silty to sandy, lignite common in basal clays. |
| | | | | | |

A-15
Drill-hole locality and identification number

Structural Cross-Section of the Soso 7.5-Minute Geologic Quadrangle

