

MISSISSIPPI DEPARTMENT OF ENVIRONMENTAL QUALITY
OFFICE OF GEOLOGY
OPEN-FILE REPORT 262

GEOLOGIC MAP of the CRYSTAL SPRINGS QUADRANGLE

Copeiah County, Mississippi

Geology by James E. Starnes, RPG


2013

DESCRIPTION OF MAP UNITS

QUATERNARY	HOLOCENE	Qal	ALLUVIUM Sand, flood plain sands, silts, and gravels.
		Qtc	CITRONELLE FORMATION Sand, yellow, orange, purple, red, pink, fine- to coarse-grained, predominantly quartzose, cross-bedded to massive, graveliferous, pea to cobble size, predominantly chert with lesser amounts of vein quartz, metaquartzite, agate, and sandstone; clay, pink to white, generally occurring as discontinuous lenses and as rip-up clasts, clasts. Conglomeratic ironstone ledges are common in the graveliferous sands at the base of the formation. The base of the Citronelle Formation is unconformable and occurs around 400 feet in elevation.
TERTIARY	MIOCENE	Tha	HATTIESBURG FORMATION Sand, gray, pale yellow to white, often indurated at surface, fine- to coarse-grained, cross-bedded to massive, predominantly quartzose with lesser amounts of chert, metaquartzite, and mica, silicified wood common. Clay, green, gray, brown, weathers white to brown, silty to sandy, locally lignitic, white opal nodules common.
		Tca	CATAHOULA FORMATION Sand, gray, pale yellow to white, fine to coarse-grained, cross-bedded to massive with rare laminar pea gravels, often indurated to orthoquartzitic at surface, predominantly quartzose with lesser amounts of chert, metaquartzite, and mica, silicified wood common; Clay, green, gray, brown, weathers white to brown, silty to sandy, can contain common opal and barite nodules, lignite common in basal clays.
		E-47	Drill-hole locality and identification number



GEOLOGIC MAP
CRYSTAL SPRINGS QUADRANGLE
Copeiah County, Mississippi



Copyright © 2013 Mississippi Department of Environmental Quality, Office of Geology

Geology field checked in 2012 using the 1963, U.S. Geological Survey 7.5-minute topographic quadrangle, Universal Transverse Mercator projection, 1927 North American datum, contour interval 10 feet, Universal Transverse Mercator projection, 1983 North American datum, GRS80 spheroid, 1000-meter Universal Transverse Mercator 1983 datum grid ticks, zone 15, shown in red, January 2013, magnetic north declination in quadrangle center is 0°22' west of true north, changing by 0°7' west per year.

Sources: The base map is derived from the Digital 2012 USTOPO; contour Mylar separate of the USGS 1963 topographic quadrangle; railroad features from the Federal Railroad Administration (FRA), 2002 edition, 1:100,000 scale; Public Land Survey System from Mississippi Automated Resource Information System (MARIS), 1:24,000 scale. Declination, National Oceanic and Atmospheric Administration (NOAA).

Geographic Information System by Daniel W. Morse, MDEQ does not warrant the accuracy or completeness of the source data. Geologic maps are only a guide to current understanding and do not eliminate the need for detailed investigations of specific sites for specific purposes.

This map was produced by the Mississippi Office of Geology in cooperation with the United States Geological Survey, National Geologic Mapping Program, under STATEMAP grant #G12AC20232.

Structural Cross-Section of the Crystal Springs 7.5-Minute Geologic Quadrangle

