



**GEOLOGIC MAP
OF THE
TISHOMINGO QUADRANGLE
MISSISSIPPI-ALABAMA
(INCLUDING MISSISSIPPI PORTION OF THE
BISHOP QUADRANGLE)**

Geology by Robert K. Merrill
1988

Digitized by Jonathan R. Leard
2020



Quadrangle map locations.
Mapped portions are shaded.



DESCRIPTION OF MAP UNITS

UNIT	DESCRIPTION
QUATERNARY	
Qal	ALLUVIUM Sand, medium- to brownish-gray, very fine- to very coarse-grained, subangular to subrounded quartz, silty, clayey; commonly contains organic matter; chert and quartzite pebbles common at base.
Qtl	LOW ELEVATION TERRACE DEPOSITS Sand, light-gray to dark reddish-brown, very fine- to very coarse-grained, subangular to subrounded quartz, silty, clayey; lower portions contain layers and lenses of flattened quartzite and quartz pebbles interspersed with rounded chert pebbles; iron staining common on pebbles. Distributed adjacent to present stream courses, at and above flood plain elevation.
Qth	HIGH ELEVATION TERRACE DEPOSITS Gravel, moderate reddish- to dark yellowish-brown, very fine- to very coarse-grained, flattened quartzite pebbles; iron staining common on outer surfaces; beds and lenses of sand, silt, and clay occur frequently in upper portions. Irregular bedding, occasional cross-bedding; ironstone cementation common. Mainly occur at elevations above 600 feet. Erosional contact at base.
CRETACEOUS	
SELMA GROUP	
Kc	COFFEE FORMATION Sand, light- to medium-gray, very fine- to medium-grained, subangular quartz, glauconitic, micaceous; frequently interbedded with silt, light- to medium-gray, clayey; thinly bedded with occasional intervals of irregular- to massive-bedded sand; occasional lenses and stringers of small chert gravel at base. Frequent thin ironstone beds; weathers to shades of reddish-brown. Unconformity at base.
EUTAW GROUP	
Ket	EUTAW FORMATION TOMBIGBEE SAND MEMBER Sand, medium light- to olive-gray, very fine- to medium-grained, subangular to subrounded quartz, well sorted, massive-bedded, glauconitic, micaceous, silty, clayey; weathers to various shades of reddish-brown. Frequent occurrence of ferruginous cemented sand molds of <i>Callianassa</i> sp. burrows.
Ke	LOWER EUTAW MEMBER Sand, medium- to olive-gray, fine- to medium-grained, subangular to subrounded quartz, glauconitic, micaceous, horizontal- and cross-bedded; commonly thinly interbedded and interlaminated with clay, medium-gray, locally carbonaceous; isolated occurrences of petrified wood in lower portions. Weathers to various shades of reddish-brown. Frequent occurrence of ferruginous cemented sand molds of <i>Callianassa</i> sp. burrows. Unconformity at base.
Kmc	McSHAN FORMATION Sand, pale yellowish-brown to very light-gray, very fine- to fine-grained, well sorted, subangular quartz, glauconitic, micaceous, silty; thinly interbedded and interlaminated with silt, light-gray to grayish orange-pink, micaceous, clayey. Horizontal- and ripple-laminated; frequent zones of massive- to cross-bedded, fine- to coarse-grained sand; frequent chert pebble lenses and stringers. Weathers to various shades of reddish-brown; yellowish-gray; local occurrences of ferruginous cemented sand molds of <i>Callianassa</i> sp. burrows; common occurrence of petrified wood; occasional occurrence of carbonaceous clays, dark-gray, micaceous, containing carbonized wood fragments. Unconformity at base.
MISSISSIPPIAN	
TUSCALOOSA GROUP	
Kt	TUSCALOOSA GROUP (UNDIFFERENTIATED) Gravel, chert, white to dark-gray, very well rounded; frequent silt and clay matrix; sand, light- to moderate reddish-brown, very fine- to very coarse-grained, subrounded to angular quartz and chert grains, poorly sorted, with frequent gravel lenses and stringers; clay, white- to medium-gray with occasional occurrences of carbonaceous dark-gray clays; zones of multi-colored chert gravel; isolated occurrences of quartzite; frequent well-cemented chert pebble conglomeratic zones. Laterally traceable silt and clay intervals occur most frequently in uppermost and lowermost intervals. Unconformity at base.
CHESTERIAN SERIES	
Mh	HARTSELLE FORMATION Sandstone, light-gray to light brownish-gray, fine- to medium-grained, well cemented quartz arenite, thin- to massive-bedded; contains thin intervals of thinly bedded and laminated siltstone and shale, medium- to dark-gray; local ferruginous staining.
Mpm	PRIDE MOUNTAIN FORMATION Shale, olive- to dark-gray, calcareous sandy, limestone, light- to brownish-gray, thin-bedded grainstones, wackestones, and mudstones, fossiliferous, occasionally oolitic, sandy, silty; sandstone, very light- to brownish-gray, thin- to massive-bedded, fine- to medium-grained, sparingly fossiliferous. Unconformity at base.
IOWA GROUP	
Ml	TUSCUMBIA FORMATION Limestone, light- to dark bluish-gray, fossiliferous, bioclastic grainstone, wackestone, and mudstone, thin- to thick-bedded, occasionally massive-bedded; some calcareous shale interbeds. Lowermost strata contain beds of chert, very dark-gray to black; uppermost strata contain grainstone, very light-gray, cross-bedded; local occurrences of nodular chert.

Base map prepared from Tishomingo Quadrangle, Tennessee Valley Authority-United States Geological Survey, 1950, photorevised in 1967. 1927 North American datum.

