

GEOLOGIC MAP of the MISSISSIPPI GULF COAST

Jackson, Harrison, and Hancock Counties

Open-File Report 355

2025

Geology by

Jonathan R. Leard, RPG, James E. Starnes, RPG, and Timothy J. Palmer, RPG




Mississippi Department of Environmental Quality
Mississippi Office of Geology - Surface Mapping Division
Mississippi Geological Survey
700 North State Street
Jackson, Mississippi 39225

Copyright © 2025 Mississippi Department of Environmental Quality, Office of Geology

This geologic map was funded by the State of Mississippi and the United States Geological Survey's National Cooperative Geologic Mapping Program.

Correlation of Map Units



Descriptions of Map Units

- Barrier Islands (Holocene)**
Sand, transported by longshore currents. Winnowed, laminated, with occurrences of heavy minerals. See Bulletin 93 for more information on heavy minerals present.
- Alluvium Fans (Holocene to Pleistocene)**
Sand, yellow to brownish-white, fine to coarse-grained, subrounded to rounded, predominantly quartzose; silt to clayey; humus lenses common; silicified wood common.
- Dunes (Holocene to Pleistocene)**
Well sorted, medium- to coarse-grained frosted quartz sand deposited by wind action with a gradual windward slope.
- Alluvium (Holocene to Pleistocene)**
Floodplain sand, silt, gravel, and clay. The Pascagoula River and Pearl River are major coastal streams with alluvial thicknesses exceeding 60 feet.
- Cut-off Channels (Holocene to Pleistocene)**
Carbonaceous clay, silt to lignitic, brown to black; includes oxbow lakes and clay plugs of abandoned meander courses.

- Coastal Terraces (Pleistocene)**
Sand, orange to tan, fine to coarse-grained, predominantly quartzose, cross-bedded to massive; clay, kaolinitic, pink to white, generally occurring as discontinuous lenses. Ferruginous sandstone and pyroclastic common at the basal contact with the underlying Graham Ferry Formation.
- Panlico**
Approximate terrace geomorphic surface at 25 feet msl.
- Big Ridge**
Approximate terrace geomorphic surface at 50 feet msl.
- Good Hope**
Approximate terrace geomorphic surface at 100 feet msl.

- River Terraces (Pleistocene)**
Fluvial fining-upward sequence of tan to khaki floodplain silts; sand, orange to tan, fine- to coarse-grained, predominantly quartzose, cross-bedded to massive; gravel, pea-sized, predominantly leached to chalky brown, gray, and white chert and milky quartz; clay, kaolinitic, pink to white, generally occurring as discontinuous lenses. Ferruginous sandstone and pyroclastic common at the basal contact with the underlying Graham Ferry and Pascagoula formations.
- Wade**
Approximate terrace geomorphic surface at 50 feet msl.
- Big Point**
Approximate terrace geomorphic surface at 70 feet msl.
- Hurley**
Approximate terrace geomorphic surface at 100 feet msl.
- Hurleston**
Approximate terrace geomorphic surface at 130 feet msl.
- Moveila**
Approximate terrace geomorphic surface at 150 feet msl.

- Graham Ferry Formation (Pliocene)**
Sand, dark greenish-gray, yellow to tan, micaceous and glauconitic (exclusively in the fine-grained sands), fine- to coarse-grained, predominantly quartzose, cross-bedded to massive; laminar to thinly bedded quartz pea gravels in the coarser fraction. Weathers to orange, purple, red, pink, with reddish-brown pebbly ironstone residuum. Clay, green, gray, brown; weathers mottled purple to pink and white to reddish-brown, silty to sandy, locally lignitic. The unconformable contact between the Pascagoula and Graham Ferry formations is projected using subsurface mapping of the Glendon Limestone and the thickness of the section between the Glendon Limestone (2450 msl) and the type locality of the Graham Ferry Formation on the west bank of the Pascagoula River at Wade to Vanceburg Road Sec. 37 T. 58 R. 7N.

- Pascagoula Formation (Miocene)**
Shallow marine to intertidal deposits, containing the marker fossil, *Rangia johnsoni*. Clay, green, gray, brown, and white; locally lignitic, locally calcareous and fossiliferous. Weathers mottled purple to pink and white to reddish-brown, silty to fine-sandy. Sand, dark greenish-gray and glauconitic, micaceous, locally lignitic, fine- to coarse-grained, predominantly quartzose; silicified wood is common.

